

University of Groningen

SecA supports a constant rate of preprotein translocation

Tomkiewicz, D; Nouwen, N; van Leeuwen, R; Tans, S; Driessen, AJM

Published in:
The Journal of Biological Chemistry

DOI:
[10.1074/jbc.M600205200](https://doi.org/10.1074/jbc.M600205200)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tomkiewicz, D., Nouwen, N., van Leeuwen, R., Tans, S., & Driessen, AJM. (2006). SecA supports a constant rate of preprotein translocation. *The Journal of Biological Chemistry*, 281(23), 15709-15713. <https://doi.org/10.1074/jbc.M600205200>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUPPLEMENTARY MATERIALS

MATERIALS AND METHODS

Effect of the protein concentration on the initial rate of translocation

To analyse the effect of the preprotein concentration on the initial rate of translocation, an *in vitro* translocation assay was performed with different concentrations of preprotein (0.5 – 0.01 μM).

FIGURE LEGEND

Fig. S1. Initial rate of translocation rate as function of the preprotein concentration. Translocation of proOmpA-P1 and P8 into IMV's derived from strain NN100 in the absence of PMF, was performed as described in Materials and Methods.




Fig. S1. Tomkiewicz *et al.*